


Derivatives of Inverse Functions



The slopes are reciprocal: $\left. \frac{df^{-1}}{dx} \right|_{f(a)} = \frac{1}{\left. \frac{df}{dx} \right|_a}$

Figure 3.52 The graphs of a function and its inverse. Notice that the tangent lines have reciprocal slopes.

THEOREM 3 Derivatives of Inverse Functions

If f is differentiable at every point of an interval I and df/dx is never zero on I , then f has an inverse and f^{-1} is differentiable at every point of the interval $f(I)$.


EXPLORATION 1 Finding a Derivative on an Inverse Graph Geometrically

Let $f(x) = x^5 + 2x - 1$. Since the point $(1, 2)$ is on the graph of f , it follows that the point $(2, 1)$ is on the graph of f^{-1} . Can you find


$$\frac{df^{-1}}{dx}(2),$$

the value of df^{-1}/dx at 2, without knowing a formula for f^{-1} ?

1. Graph $f(x) = x^5 + 2x - 1$. A function must be one-to-one to have an inverse function. Is this function one-to-one?
2. Find $f'(x)$. How could this derivative help you to conclude that f has an inverse?
3. Reflect the graph of f across the line $y = x$ to obtain a graph of f^{-1} .
4. Sketch the tangent line to the graph of f^{-1} at the point $(2, 1)$. Call it L .
5. Reflect the line L across the line $y = x$. At what point is the reflection of L tangent to the graph of f ?
6. What is the slope of the reflection of L ?
7. What is the slope of L ?
8. What is $\frac{df^{-1}}{dx}(2)$?



Derivative of the Arcsine



$$y = \sin^{-1} x$$

$$\cos^2 y = 1 - \sin^2 y$$

Inverse Trig Function Derivatives Activity

$$\frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}.$$

If u is a differentiable function of x with $|u| < 1$, we apply the Chain Rule to get

$$\frac{d}{dx} \sin^{-1} u = \frac{1}{\sqrt{1-u^2}} \frac{du}{dx}, \quad |u| < 1.$$

EXAMPLE 1 Applying the Formula

$$\frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\sin^{-1} x^2)$$

$$= \frac{1}{\sqrt{1-(x^2)^2}} \cdot 2x$$

$$= \frac{2x}{\sqrt{1-x^4}}$$

EXAMPLE 2 A Moving Particle

A particle moves along the x -axis so that its position at any time $t \geq 0$ is $x(t) = \tan^{-1} \sqrt{t}$. What is the velocity of the particle when $t = 16$?

$$\frac{d}{dx} \tan^{-1} x = \frac{1}{1+x^2}$$

$$\begin{aligned} \frac{d}{dt} \tan^{-1} \sqrt{t} &= \frac{d}{dt} \tan^{-1} t^{1/2} \\ &= \frac{1}{1+(t^{1/2})^2} \cdot \frac{1}{2\sqrt{t}} \\ &= \frac{1}{2\sqrt{t}(1+t)} \\ &= \frac{1}{136} \end{aligned}$$

EXAMPLE 3 Using the Form

$$\frac{d}{dx} \sec^{-1}(5x^4) =$$

$$\sec^{-1} x = \frac{1}{|x| \sqrt{x^2 - 1}}$$

$$x = \sec y$$

$$\tan^2 y = \sec^2 y - 1$$

$$\tan y = \pm \sqrt{\sec^2 y - 1}$$

$$\sec y \tan y = \frac{dy}{dx}$$

$$\rightarrow \frac{1}{|5x^4| \sqrt{(5x^4)^2 - 1}} \cdot 20x^3$$

$$= \frac{20x^3}{5x^4 \sqrt{25x^8 - 1}}$$

$$= \frac{4}{x \sqrt{25x^8 - 1}}$$

Inverse Function-Inverse Cofunction Identities

$$\cos^{-1} x = \pi/2 - \sin^{-1} x$$

$$\cot^{-1} x = \pi/2 - \tan^{-1} x$$

$$\csc^{-1} x = \pi/2 - \sec^{-1} x$$

Use these identities to come up with the derivatives of these 3 inverse functions.

$$\frac{d}{dx} \cos^{-1} x = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \cot^{-1} x = \frac{-1}{1+x^2}$$

$$\frac{d}{dx} \csc^{-1} x = \frac{-1}{|x|\sqrt{x^2-1}}$$

Calculator Conversion Identities

$$\sec^{-1} x = \cos^{-1} (1/x)$$

$$\cot^{-1} x = \pi/2 - \tan^{-1} x$$

$$\csc^{-1} x = \sin^{-1} (1/x)$$

Notice that we do not use $\tan^{-1} (1/x)$ as an identity for $\cot^{-1} x$. A glance at the graphs of $y = \tan^{-1} (1/x)$ and $y = \pi/2 - \tan^{-1} x$ reveals the problem (Figure 3.55).




Figure 3.55 The graphs of (a) $y = \tan^{-1} (1/x)$ and (b) $y = \pi/2 - \tan^{-1} x$. The graph in (b) is the same as the graph of $y = \cot^{-1} x$.

EXAMPLE 4 A Tangent Line to the Arccotangent Curve

Find an equation for the line tangent to the graph of $y = \cot^{-1} x$ at $x = -1$.